

EAST Search History

Ref #	Hits	Search Query	DBs	Default Operator	Plurals	Time Stamp
L1	1032	((546/298) or (514/350)).CCLS.	US-PGPUB; USPAT; USOCR; EPO; JPO; DERWENT	OR	OFF	2007/01/22 12:09
L2	2238542	"2006".py. or "2007".py.	US-PGPUB; USPAT; USOCR; EPO; JPO; DERWENT	OR	ON	2007/01/22 12:09
L3	100	1 and 2	US-PGPUB; USPAT; USOCR; EPO; JPO; DERWENT	OR	ON	2007/01/22 12:09

Dare to compare?
Find out how your hospital compares with others



NPL Search Notes

[ACP Medicine] [MedGenMed eJournal]

Latest | News | CME | Conferences | Resource Centers | Patient Ed. | Journals & Reference | Experts & Viewpoints

Search Medscape, MEDLINE and Drug Reference

Search

Newsletters | Log In | Register

Is your patient's
drug regimen
safe?

Now it's easy
to find out.

SEARCH

Medscape
 MEDLINE
 Drug Reference
[Advanced Search](#)

NEW in
Drug Reference:
Multi-Drug
Interaction
Checker
from Medscape.com

Check for
drug interactions
[click here](#)

Article Tools

Printer-Friendly Email This Download This View Online

Alteration of lacto-series glycolipid glycosyltransferase activities in human colonic adenocarcinoma DLD-1 cells after culture in N,N-dimethylformamide-containing medium.

J Cell Biochem. 1990; 44(2):93-105 (ISSN: 0730-2312)

Holmes EH; Greene TG
Pacific Northwest Research Foundation, Seattle, Washington
98122.

* Human colonic adenocarcinoma DLD-1 cells were grown under conditions which induce characteristics of differentiated cells using medium containing 0.8% N,N-dimethylformamide in order to study alterations in glycosphingolipid glycosyltransferase activities during this process. Analysis of biosynthetic reactions involved in lacto-series antigen synthesis revealed no changes in the specific activities of either beta 1----4galactosyltransferase or alpha 1----3/4fucosyltransferase with N,N-dimethylformamide treatment. However, a dramatic decrease of from 14- to 20-fold in the beta 1----3N-acetylglucosaminyltransferase activity was observed in the treated cells. This enzyme catalyzes the rate-limiting step in lacto-series core chain synthesis. This is consistent with the pattern of regulation of lacto-series antigen expression found to occur during oncogenesis in human colonic mucosa (Holmes EH, Hakomori S, Ostrander GK: J Biol Chem 262:15649, 1987). Total glycolipids from untreated and N,N-dimethylformamide-treated cells were isolated and subjected to TLC immunostain analysis and solid phase radioimmunoassay with a series of monoclonal antibodies specific for lacto-series-based carbohydrate antigens. A decrease of about 2-fold or less in the quantity of lacto-series antigens was observed as a consequence of N,N-dimethylformamide treatment in both neutral glycolipid and ganglioside fractions. The results suggest that only very low levels of beta 1----3N-acetylglucosaminyltransferase activity are required for the steady state expression of significant levels of lacto-series based glycolipids and that modulation of its activity levels by N,N-dimethylformamide treatment in DLD-1 cells represents a convenient in vitro system for studying aspects of regulation of lacto-series antigen expression.

► Subject Headings

PreMedline Identifier: 2250046

Information from Industry

Antibiotic Treatment for Many Outpatient and Inpatient Bacterial Infections

Learn more about high-dose, short-course therapy, including PK/PD parameters, clinical benefits, and doctor and patient benefits.

Medscape Newsletters

Sign Up To Receive Medscape Best Evidence

Key journal articles ranked for newsworthiness and clinical relevance in each specialty, linked to Medline abstracts.

Search Medscape, MEDLINE and Drug Reference

Search